

# Relationship between control and the intermediate term outcome of anorexia nervosa in Hong Kong

Sing Lee, Y.Y. Lydia Chan, Kathleen Kwok, L.K. George Hsu

Australian, New Zealand Journal Of Psychiatry, 2005, March 39(3) 141-145

<https://www.ncbi.nlm.nih.gov/pubmed/15701062>

put abstract and material below

Department of Psychiatry and Hong Kong Eating Disorders Center, The Chinese University of Hong Kong, 7 A, Block E, Staff Quarters, Prince of Wales Hospital, Shatin, Hong Kong. Email: [singlee@cuhk.edu.hk](mailto:singlee@cuhk.edu.hk)

Y.Y. Lydia Chan, former Research Assistant

Department of Psychiatry, The Chinese University of Hong Kong, Hong Kong

Kathleen Kwok, Clinical Psychologist

Hong Kong Eating Disorders Center, The Chinese University of Hong Kong, Hong Kong

L.K. George Hsu, Professor

Department of Psychiatry, Tufts University School of Medicine, MA, USA

Sing Lee, Professor (Correspondence)

## Abstract

**Objective:** To examine the relationship between control and the intermediate term outcome of Chinese patients with anorexia nervosa.

**Method:** 88 patients who fulfilled the DSM-III-R criteria for typical (fat phobic, n = 63) and atypical (non-fat phobic, n = 25) anorexia nervosa were contacted 9 years after the onset of illness. They completed the Morgan-Russell Outcome Assessment Schedule, Shapiro Control Inventory, and Eating Disorder Inventory-I.

**Results:** 62.2%, 32.4% and 5.4% of patients had good, intermediate, and poor outcome. Outcome was positively correlated with the overall general and specific sense of control, positive sense of control, and negatively with negative sense of control. Patients with good outcome exhibited the least negative modes of control and had the least desire for control. Typical and atypical patients showed similar but not identical control profiles.

**Conclusion:** Healthier control was associated with better outcome of anorexia nervosa but their causal relationship was far from clear because of the cross-sectional nature of the present study. The Shapiro Control Inventory is a potentially useful instrument for studying control in eating disorders.

\* \* \*

